


Syphilis



Approved 04/03/2002
 Copyright © April, 2002
 UNC School of Medicine
 at Chapel Hill

Syphilis References

at Chapel Hill. They cannot be reproduced in whole or part without the *expressed* permission of the school.

Copyright © April 2002 UNC School of Medicine at Chapel Hill

1 . Centers for Disease Control and Prevention. 1998 Guidelines for treatment of sexually transmitted diseases. MMWR 1998; 47(no.RR-1):[29] *The VDRL and RPR are equally valid.*

2. Centers for Disease Control and Prevention. 1998 Guidelines for treatment of sexually transmitted diseases. MMWR 1998; 47(no.RR-1):[31] *Patients who have primary or secondary syphilis should be treated with Benzathine penicillin G 2.4 million units IM in a single dose.*

3 . Centers for Disease Control and Prevention. 1998 Guidelines for treatment of sexually transmitted diseases. MMWR 1998; 47(no.RR-1):[34] *The following regimes are recommended for non allergic patients who have normal CSF examination (if performed) with Benzathine penicillin G 7.2 million units total, administered as three doses of 2.4 million units IM each at 1-week intervals.*

4 . Centers for Disease Control and Prevention. 1998 Guidelines for treatment of sexually transmitted diseases. MMWR 1998; 47(no.RR-1):[41] *Coordinated prenatal care and treatment follow up are important, and syphilis case management may help facilitate prenatal enrollment. Serologic titers should be repeated in the third trimester and at delivery. Serological titers may be checked monthly in women at high risk for re-infection or in geographic areas in which the prevalence of syphilis is high.*

NOTIFICATION TO USERS

These algorithms are designed to assist the primary care provider in the clinical management of a variety of problems that occur in pregnancy. They should not be interpreted as *standard of care* but instead represent *guidelines* for the management of these patients. Variation in practice should be taken into account such factors as characteristics of the individual patient, health resources, and regional experience with diagnostic and therapeutic modalities. The algorithms remain the intellectual property of the University of North Carolina School of Medicine

