


UNC Center for Maternal & Infant Health

Progesterone for prevention of preterm birth



Use of progesterone or cerclage in multiple gestation, as well as the utility of US cervical screening in multiple gestation remains unproven. Progesterone has not been demonstrated to prevent preterm birth in unselected twins/triplets.

References

- 1. Campbell S. Universal cervical length screening and vaginal progesterone prevents early preterm births, reduces neonatal morbidity and is cost saving: doing nothing is no longer and option. Ultrasound Obstet Gynecol. 38 (1), 1-9 (2011).
- 2. Meis PJ. Prevention of recurrent preterm delivery by 17 alpha hydroxyprogesterone caproate. N Engl J Med 2003; 348; 2379-85.
- 3. ACOG Committee Opinion No. 387 Pharmaceutical Compounding, November 2007, http://www.acog.org/partners/greenjournal/co387.pdf .
- 4. http://www.acog.org/from home/misc/20111013MakenaLtr.pdf.
- 5. http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm249025.htm.
- 6. Fonseca, EB. Progesterone and the risk of preterm birth among women with a short cervix. N Engl J Med 2007 Aug, 2;357(5): 462-9.
- 7. Hassan SS. Vaginal progesterone reduces the rate of preterm birth in women with a sonographic short cervix: a multicenter, randomized, double-blind, placebo-controlled trial. Ultrasound Obstet Gynecol. 2011 Jul;38(1):18-31.
- 8. Iams JD, Berghella V Care for women with prior preterm birth. Am J Obstet Gynecol 2010.
- 9. Romero, R, et al. vaginal Progesterone in women with asymptomatic sonographic short cervix in the midtrimester decreases preterm delivery and neonatal morbidity: A systematic review and meta-analysis of individual patient level data. Am J Obstet Gynecol. 201.

Revised January 17, 2012.

Notification to Users

These algorithms are designed to assist the primary care provider in the clinical management of a variety of problems that occur during pregnancy. They should not be interpreted as a standard of care, but instead represent guidelines for management. Variation in practices should take into account such factors as characteristics of the individual patient, health resources, and regional experience with diagnostic and therapeutic modalities.

The algorithms remain the intellectual property of the University of North Carolina at Chapel Hill School of Medicine. They cannot be reproduced in whole or in part without the expressed written permission of the school.