

Newborn Critical Care Center (NCCC) Clinical Guidelines

Premedication for Non-Emergent Neonatal Intubation

INTRODUCTION

Endotracheal intubation should be performed in a controlled environment with appropriate preparation whenever the patient's status permits to prevent traumatic injury to the airway/oral tissues. Intubation *without* premedication has been associated with significant side effects including pain, hypoxia and hemodynamic changes.

Benefits of premedication for non-emergent intubation include:

- Decreased pain
- Fewer number of attempts to complete the intubation, resulting in decreased airway trauma
- Higher success rates for inexperienced personnel
- Decreased hypoxia and hemodynamic effects (bradycardia, increased intracranial pressure, risk of IVH, systemic and pulmonary hypertension)

The effects of premedication are maximized when the full complement of drugs are used together.

An intubation may be considered non-emergent when the following criteria are met:

- The intubation is not occurring in the delivery room
- The infant has intravenous access, or is stable enough to delay intubation until intravenous access is obtained
- The infant is stable enough to delay intubation until medications can be obtained and administered

PROCEDURE

1. **Order medications:** In EPIC search for the Neonatal Premedication Order Panel, which can also be accessed through the "Neonatal Procedure Focused" order set, then choose "Neonatal Intubation Premedications."
2. Medication doses should be calculated and checked by two nurses.
3. Provider, RN and RT at bedside should perform timeout prior to administration of

DRUGS <i>(GIVE IN THE ORDER LISTED)</i>	DOSE	ROUTE	ONSET (MINUTES)	DURATION
Atropine	0.02 mg/kg	IV push	1-2 minutes	2-4 hours
Fentanyl	2 mcg/kg	IV over 5 min	1-2 minutes	30-60 minutes
Rocuronium <i>* Paralytics should never be administered alone</i>	0.5 mg/kg	IV push	0.2 - 2 minutes <i>* May be repeated if not effective in 1.5 – 2 minutes</i>	30-40 minutes

4. Medications **should be given in the physical presence of a medical provider.**
5. Intubation should be attempted when cessation of spontaneous movement is noted and laxity of the jaw is determined. (Refer to [NCCC Intubation Guidelines](#))
6. Respiratory therapy is expected to be present for the entire procedure and additional medical providers should be available to attempt intubation if necessary.

POST-PROCEDURE

After the ETT has been secured, the patient should continue to be closely monitored and vital signs recorded. The provider should ensure that there is sufficient respiratory support immediately after intubation to achieve the desired level of minute ventilation. As the effects of the medications diminish, it may be appropriate to reduce this level of respiratory support.

References:

1. Allen, KA. Premedication for Neonatal Intubation: Which Medications are recommended and why? *Advances in Neonatal Care* (2012) 12(2):107-111.
2. Foglia E, Ades A, Sawyer T et al. Neonatal intubation practice and outcomes: An international registry study. *Pediatrics*. 2019 Jan; 143(1): e20180902.
3. Hatch, L.D., Grubb, P. H., Lea, A.S., Walsh, W.F., Markham, M.H., Maynard, P. O. Whitney, G.M. Stark, A.R., & Ely, W. Interventions to Improve Patient Safety During Intubation in the Neonatal Intensive Care Unit. *Pediatrics* (2016) Oct 138(4):
4. Le, CN, Garey DM, Leone, TA, Goodmar, JK, Rich, W. and Finer, N. [Impact of premedication on neonatal intubations by pediatric and neonatal trainees](#). *Journal of Perinatology* (2014) 34, 458–460
5. Muniraman, HK., Yaari J., Hand, I. [Premedication Use Before Nonemergent Intubation in the Newborn Infant](#). *Am J Perinatol* 2015; 32:821–824.
6. North Carolina Board of Nursing. (2022). *Rapid sequence intubation (RSI)* [Position statement]. Retrieved from <https://www.ncbon.com/myfiles/downloads/position-statements-decision-trees/rapid-sequence-intubation.pdf>
7. O'Connor, T. L. Premedication for nonemergent neonatal intubation: A systematic review. *Journal of Perinatal Neonatal Nursing* (2022); 36 (3), 284-296.
8. Ozawa, Y., et al. Premedication with neuromuscular blockade and sedation during neonatal intubation is associated with fewer adverse events. *Journal of Perinatology*. (2019). 39:848-856
9. Shay, R, Weikel, B.W., et al. Standardizing premedication for non-emergent neonatal tracheal intubations improves compliance and patient outcomes. *Journal of Perinatology* (2021). 42: 132-138.